

I.P.A Banned Substance List

The I.P.A. was established with a separate drug tested division. The Amateur division is a strict drug tested division. Lifters in the Amateur division will be tested by urinalysis for the presence of prohibited anabolic steroids. With the continued growth of the I.P.A., it is the Association's plan to test more lifters.

Below is a list of *anabolic steroids* and recommended I.P.A. Laboratories that can be used for testing

Bolasterone
Boldenome (Equipoise)
4-Chlorotestosterone(Clostebol)
Clenbuterol (Siropent)
Danazol
Dihydrotestosterone (Stanolone)
Dehydrochloromethyltestosterone (Oral Turinabol)
Dromostanolone (Masteron)
Epitestosterone (if above 200 ng/mL)
Formebolone (Esiclone)
Fluoxymesterone (Halotestin)
Furazabol
Mesterolone (Proviron)
Methandrostenolone (Methandienone, Dianabol)

Methandriol
Methenolone (Primobolan)
Methyltestosterone
Mibolerone
Nandrolone (Deca-Durabolin)
Norethandrolone/Ethylestrenol
Oxandrolone (Anavar)
Oxymesterone
Oxymetholone (Anadrol)
Stanozolol (Winstrol)
Stenbolone
Testosterone (T/epiT ratio*)
Trenbolone (Finajet, Parabolan)
... *and other related compounds*
Probenecid (masking agent)

*specimen is considered positive when Testosterone/Epitestosterone ratio is greater than six (6).